

575

Yellow Peel · Leave-On

Melasma · PIH · Photoaging · Dull / Uneven Tone
Fitzpatrick II–VI — gentle

RG-2 Retinol / Glycolic

getprodermic.com/protocols/575-yellow-peel-leave-on CLICK TO OPEN THE PRODUCT PAGE

30ML · ~20 APPLICATIONS

WEAR MODEL — APPLIED IN CLINIC, RINSED AT HOME

No frost expected — mild warmth only. Patient rinses off at home after 8–10 hours.

10 HR
MAX WEAR

TIME	PROCEDURE	STOP — NEUTRALIZE
2 MIN	Prep Cleanse + degrease (PREP.01), air-dry. First session: patch test post-auricular, observe 10 min.	Abort on any of these, at any minute. NEUT.01 first, cold water second.
1 MIN	Apply Thin, even layer by fan brush. Avoid eyes, lips, nostrils. No layering.	Escalating burning or pain
8–10 HR	Wear Patient leaves with the peel on. Wear 8–10 hours (or overnight per your protocol), then rinse.	Deep-red erythema beyond the expected uniform pink
HOME	Rinse Patient rinses at home with water. Brief them on NEUT.01-style cool-water relief at first sign of over-reaction.	Edema in the treated field
—	Finish Balm (REST.01) + SPF 50+ after the home rinse. Hand over the aftercare card.	Urticaria or wheals
		Patient cannot tolerate it
		PROGRAM 4–6 sessions per course 25–30 days between sessions Allow 25–45 days between any peels , by strength and potency. Prime higher phototypes: 2 weeks topical retinoid.

DO NOT TREAT IF Re-verify every visit, not only the first.

- ✗ Active dermatitis, eczema or infection
 - ✗ Topical retinoid within 72 hours
 - ✗ Allergy to any listed active
 - ✗ Sunburned / recent sun exposure
- ✗ Pregnancy or breastfeeding
 - ✗ Broken skin / compromised barrier
 - ✗ Laser / IPL within 2 weeks
- ✗ Oral isotretinoin within 6 months
 - ✗ Keloid or hypertrophic scar history
 - ✗ Active herpes simplex

PATIENT AFTERCARE

DAY 0	DAY 1–2	DAY 3–5	DAY 5–7	ONGOING
Keep clean. Balm only.	Moisturize 3–4x daily.	Peeling. Never pick.	Recovery. Makeup once shed.	Daily SPF 50+ . Actives: day 7.