

G70

Glycolic Peel

Anti-Aging · Resurfacing · Acne · Brightening
Fitzpatrick I-III — caution IV

GA-70 Glycolic

getprodermic.com/protocols/g70-glycolic-peel CLICK TO OPEN THE PRODUCT PAGE



30ML

ENDPOINT — TREAT TO ENDPOINT, NOT TO THE CLOCK

Uniform pink erythema → light frost. Then neutralize.

5 MIN
MAX CONTACT

CAUTION Very high potency. Hard-timed contact — set a timer at application. Mandatory neutralization with NEUT.01; never exceed 5 minutes.

TIME	PROCEDURE
2 MIN	Prep Cleanse + degrease (PREP.01), air-dry. First session: patch test post-auricular, observe 10 min.
1 MIN	Apply Thin, even layer by fan brush. Avoid eyes, lips, nostrils. No layering.
2-5 MIN	Titrate Observe continuously to endpoint. Start low on first sessions; extend only with proven tolerance. Never exceed 5 minutes.
2 MIN	Neutralize NEUT.01 — every session, every zone. Cold water as secondary. Never let it self-neutralize.
—	Finish Balm (REST.01) + SPF 50+ before the patient leaves. Hand over the aftercare card.

STOP — NEUTRALIZE

Abort on **any** of these, at any minute. NEUT.01 first, cold water second.

Escalating burning or pain

Deep-red erythema
beyond the expected uniform pink

Edema in the treated field

Urticaria or wheals

Patient cannot tolerate it

PROGRAM

4-6 sessions per course

40-45 days between sessions

Allow 25-45 days between any peels, by strength and potency.

DO NOT TREAT IF Re-verify every visit, not only the first.

- ✗ Active dermatitis, eczema or infection
 - ✗ Topical retinoid within 72 hours
 - ✗ Allergy to any listed active
 - ✗ Sunburned / recent sun exposure
- ✗ Pregnancy or breastfeeding
 - ✗ Broken skin / compromised barrier
 - ✗ Laser / IPL within 2 weeks
- ✗ Oral isotretinoin within 6 months
 - ✗ Keloid or hypertrophic scar history
 - ✗ Active herpes simplex

PATIENT AFTERCARE

DAY 0	DAY 1-2	DAY 3-5	DAY 5-7	ONGOING
Keep clean. Balm only.	Moisturize 3-4x daily.	Peeling. Never pick.	Recovery. Makeup once shed.	Daily SPF 50+ . Actives: day 7.

For professional use by trained practitioners only. Supports — does not replace — clinical judgment.
Obtain informed consent and baseline photographs. Expected downtime: 1-3 days.